

DEFORMATION QUANTISATION FOR UNSHIFTED SYMPLECTIC STRUCTURES ON DERIVED ARTIN STACKS

J.P. PRIDHAM

ABSTRACT. We construct a map from DQ algebroid quantisations of unshifted symplectic structures on a derived Artin N -stack to power series in de Rham cohomology, depending only on a choice of Levi decomposition for the Grothendieck–Teichmüller group. This gives an equivalence between even power series and certain involutive quantisations, which yield involutive curved A_∞ deformations of the dg category of perfect complexes. In particular, there is a canonical quantisation associated to every symplectic structure on such a stack, which agrees for smooth varieties with the Kontsevich–Tamarkin quantisation.

INTRODUCTION

For $n > 0$, existence of quantisations of n -shifted Poisson structures is a formality, following from the equivalence $E_{n+1} \simeq P_{n+1}$ of operads. Quantisations of positively shifted symplectic structures thus follow immediately from the equivalence in [Pri5, CPT⁺] between symplectic and non-degenerate Poisson structures. In [Pri4], quantisation for non-degenerate (-1) -shifted Poisson structures was established, and we now consider the $n = 0$ case, fleshing out the details sketched in [Pri4, §4.3].

For Poisson structures on smooth algebraic varieties, the Kontsevich–Tamarkin quantisations rely on formality of the Hochschild complex, so will not adapt to arbitrary Poisson structures on derived stacks. We show, however, that all non-degenerate Poisson structures can be quantised even when the Hochschild complex is not formal, by a similar mechanism to the quantisation of non-degenerate (-1) -shifted Poisson structures in [Pri4].

The proof in [Pri5] of the correspondence between n -shifted symplectic and non-degenerate Poisson structures relied on the existence, for all Poisson structures π , of a CDGA morphism $\mu(-, \pi)$ from the de Rham algebra to the algebra $T_\pi \widehat{\text{Pol}}(X, n)$ of shifted polyvectors with differential twisted by π . In [Pri4], this idea was extended to establish the existence of quantisations for (-1) -shifted symplectic structures, with μ being an A_∞ -morphism from the de Rham algebra to the ring of differential operators.

In order to adapt these constructions to 0-shifted symplectic structures, we replace polyvectors or differential operators with the Hochschild complex $\text{CC}_R^\bullet(X)$ of X over R . Since this has an E_2 -algebra structure, a choice w of Levi decomposition for the Grothendieck–Teichmüller group gives it a P_2 -algebra structure. Any quantisation Δ (a Maurer–Cartan element in $\text{CC}_R^\bullet(X)[[\hbar]]$) then defines a morphism $\mu_w(-, \Delta)$ from the de Rham complex $\text{DR}(X/R)$ to $\text{CC}_R^\bullet(X)[[\hbar]]$ twisted by Δ . In more detail, since $[\Delta, -]$ defines a derivation from \mathcal{O}_X to $\text{CC}_R^\bullet(X)[[\hbar]]$, it determines a map $\Omega_X^1 \rightarrow \text{CC}_R^\bullet(X)[[\hbar]][1]$ and $\mu_w(-, \Delta)$ is the resulting morphism of CDGAs.

This work was supported by the Engineering and Physical Sciences Research Council [grant number EP/I004130/2].

This gives rise to a notion of compatibility between E_1 quantisations Δ and generalised pre-symplectic structures (power series ω of elements of the de Rham complex): we say that ω and Δ are w -compatible if

$$\mu_w(\omega, \Delta) \simeq \hbar^2 \frac{\partial \Delta}{\partial \hbar}.$$

Proposition 2.16 shows that every non-degenerate affine quantisation Δ has a unique w -compatible generalised pre-symplectic structure, thus giving us a map

$$Q\mathcal{P}(A, 0)^{\text{nondeg}} \rightarrow H^2(F^2\text{DR}(A)) \times \hbar H^2(F^1\text{DR}(A)) \times \hbar^2 H^1(\text{DR}(A))[[\hbar]]$$

on the space of non-degenerate E_1 quantisations.

Moreover, we have spaces $Q\mathcal{P}(A, 0)/G^{k+1}$ consisting of E_1 quantisations of order k , by which we mean Maurer–Cartan elements in $\prod_{j \geq 2} (F_j \text{CC}_R^\bullet(X)/F_{j-k-1}) \hbar^{j-1}$, for F the good truncation filtration in the Hochschild direction. Via induction on levels of the filtration, and an analysis of the associated DGLA obstruction theory, Proposition 2.17 then shows that the resulting map

$$Q\mathcal{P}(A, 0)^{\text{nondeg}} \rightarrow (Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^2) \times \hbar^2 H^2(\text{DR}(A))[[\hbar]]$$

underlies an equivalence. Thus quantisation reduces to a first order problem.

This first order problem is resolved by introducing a notion of self-duality. In [Pri4], self-dual quantisations were defined for line bundles \mathcal{L} with an involution $\mathcal{L} \simeq \mathcal{L}^\vee$ to the Grothendieck–Verdier dual. The analogous notion in our setting is given by considering involutive associative algebras and categories. Explicitly, when X is a smooth variety, a self-dual quantisation of \mathcal{O}_X is an associative deformation

$$(\mathcal{O}_X[[\hbar]], \star_\hbar)$$

of \mathcal{O}_X with $a \star_{-\hbar} b = b \star_\hbar a$. More generally, a self-dual quantisation of X leads to a curved A_∞ -category with semilinear contravariant involution, deforming the dg category of perfect complexes on X .

Restricting to self-dual quantisations ensures that the first-order obstruction vanishes, leading to Theorem 2.20, which shows that the equivalence class of self-dual quantisations of a given non-degenerate Poisson structure is parametrised by

$$\hbar^2 H^2(\text{DR}(A))[[\hbar^2]],$$

and in particular such quantisations always exist. Global versions of these results for derived Artin N -stacks are summarised in Theorem 3.12.

The structure of the paper is as follows.

In §1 we recall the description from [Pri5] of bi-CDGAs as formal completions of derived N -stacks along derived affines, together with the complex of polyvectors $\widehat{\text{Pol}}(A, 0)$ on such objects, and the space $\mathcal{P}(A, 0)$ of Poisson structures. We then introduce a quantisation $Q\widehat{\text{Pol}}(A, 0)$ of the complex of polyvectors, defined in terms of the Hochschild complex, and introduce an involution of this complex whose fixed points give rise to self-dual quantisations.

§2 contains the technical heart of the paper. After introducing generalised pre-symplectic structures as de Rham power series, and after recalling formality quasi-isomorphisms for the E_2 operad associated to Levi decompositions w of the Grothendieck–Teichmüller group, we introduce the notion (Definition 2.12) of w -compatibility between quantisations and generalised pre-symplectic structures. The main results (Propositions 2.16, 2.17 and Theorem 2.20) then follow, establishing the

existence of quantisations of non-degenerate unshifted Poisson (and hence symplectic) structures on stacky derived affines. Proposition 2.23 shows that for suitable choices of Levi decomposition, constant power series correspond to Kontsevich–Tamarkin quantisations.

In §3, these results are translated into the fully global setting of derived Artin N -stacks (Theorem 3.12). The approach precisely mimics that of [Pri5, §§2, 3], by establishing étale functoriality in an ∞ -category setting and applying descent arguments. Proposition 3.10 shows how E_1 quantisations in our sense give rise to curved A_∞ deformations of the dg category of perfect complexes on a derived Artin N -stack.

CONTENTS

Introduction	1
1. Quantisation for stacky thickenings of derived affine schemes	3
1.1. Stacky thickenings of derived affines	3
1.2. Polyvectors	5
1.3. The Hochschild complex of a bi-CDGA	6
1.4. Quantised 0-shifted polyvectors and quantisations	7
1.5. The centre of a quantisation	10
1.6. Self-dual quantisations	11
2. Quantisations and de Rham power series	12
2.1. Generalised pre-symplectic structures	12
2.2. Formality	13
2.3. Compatible quantisations	14
2.4. The equivalences	15
2.5. Comparison with Kontsevich–Tamarkin quantisations	18
3. Quantisation for derived stacks	19
3.1. Quantised polyvectors for diagrams	19
3.2. Hypergroupoids	21
3.3. Global quantisations	22
References	24

1. QUANTISATION FOR STACKY THICKENINGS OF DERIVED AFFINE SCHEMES

1.1. Stacky thickenings of derived affines. We now recall some definitions and lemmas from [Pri5, §3], as summarised in [Pri4, §3.1]. By default, we will regard the CDGAs in derived algebraic geometry as chain complexes $\dots \xrightarrow{\delta} A_1 \xrightarrow{\delta} A_0 \xrightarrow{\delta} \dots$ rather than cochain complexes — this will enable us to distinguish easily between derived (chain) and stacky (cochain) structures.

Definition 1.1. A bi-CDGA is a chain cochain complex A_\bullet^\bullet equipped with a commutative product $A \otimes A \rightarrow A$ and unit $\mathbb{Q} \rightarrow A$. Given a chain CDGA R , a bi-CDGA over R is then a morphism $R \rightarrow A$ of bi-CDGAs. We write $DGdg\mathcal{C}Alg(R)$ for the category of bi-CDGAs over R , and $DG^+dg\mathcal{C}Alg(R)$ for the full subcategory consisting of objects A concentrated in non-negative cochain degrees.

When working with chain cochain complexes V_\bullet^\bullet , we will usually denote the chain differential by $\delta: V_j^i \rightarrow V_{j-1}^i$, and the cochain differential by $\partial: V_j^i \rightarrow V_j^{i+1}$. Readers

interested only in DM (as opposed to Artin) stacks may ignore the stacky part of the structure and consider only chain CDGAs $A_\bullet = A_\bullet^0$ throughout this section.

Definition 1.2. Say that a morphism $U \rightarrow V$ of chain cochain complexes is a levelwise quasi-isomorphism if $U^i \rightarrow V^i$ is a quasi-isomorphism for all $i \in \mathbb{Z}$. Say that a morphism of bi-CDGAs is a levelwise quasi-isomorphism if the underlying morphism of chain cochain complexes is so.

The following is [Pri5, Lemma 3.4]:

Lemma 1.3. *There is a cofibrantly generated model structure on bi-CDGAs over R in which fibrations are surjections and weak equivalences are levelwise quasi-isomorphisms.*

There is a denormalisation functor D from non-negatively graded CDGAs to cosimplicial algebras, with left adjoint D^* as in [Pri1, Definition 4.20]. Given a cosimplicial chain CDGA A , D^*A is then a bi-CDGA in non-negative cochain degrees. By [Pri5, Lemma 3.5], D^* is a left Quillen functor from the Reedy model structure on cosimplicial chain CDGAs to the model structure of Lemma 1.3.

Since DA is a pro-nilpotent extension of A^0 , when $H_{<0}(A) = 0$ we think of the simplicial hypersheaf $\mathbf{R}\mathrm{Spec} DA$ as a stacky derived thickening of the derived affine scheme $\mathbf{R}\mathrm{Spec} A^0$.

Definition 1.4. Given a chain cochain complex V , define the cochain complex $\hat{\mathrm{Tot}} V \subset \mathrm{Tot}^{\Pi} V$ by

$$(\hat{\mathrm{Tot}} V)^m := \left(\bigoplus_{i < 0} V_{i-m}^i \right) \oplus \left(\prod_{i \geq 0} V_{i-m}^i \right)$$

with differential $\partial \pm \delta$.

Definition 1.5. Given a bi-CDGA A and A -modules M, N in chain cochain complexes, we define internal Homs $\mathcal{H}om_A(M, N)$ by

$$\mathcal{H}om_A(M, N)_j^i = \mathrm{Hom}_{A_\#}(M_\#^{\#}, N_{\#[j]}^{\#[i]}),$$

with differentials $\partial f := \partial_N \circ f \pm f \circ \partial_M$ and $\delta f := \delta_N \circ f \pm f \circ \delta_M$, where $V_\#^{\#}$ denotes the bigraded vector space underlying a chain cochain complex V .

We then define the Hom complex $\underline{\hat{\mathrm{Hom}}}_A(M, N)$ by

$$\underline{\hat{\mathrm{Hom}}}_A(M, N) := \hat{\mathrm{Tot}} \mathcal{H}om_A(M, N).$$

Note that there is a multiplication $\underline{\hat{\mathrm{Hom}}}_A(M, N) \otimes \underline{\hat{\mathrm{Hom}}}_A(N, P) \rightarrow \underline{\hat{\mathrm{Hom}}}_A(M, P)$ (the same is not true for $\mathrm{Tot}^{\Pi} \mathcal{H}om_A(M, N)$ in general).

Definition 1.6. A morphism $A \rightarrow B$ in $DG^+ dg\mathrm{CAlg}(R)$ is said to be homotopy formally étale when the map

$$\{\mathrm{Tot} \sigma^{\leq q}(\mathbf{L}\Omega_A^1 \otimes_A^{\mathbf{L}} B^0)\}_q \rightarrow \{\mathrm{Tot} \sigma^{\leq q}(\mathbf{L}\Omega_B^1 \otimes_B^{\mathbf{L}} B^0)\}_q$$

on the systems of brutal cotruncations is a pro-quasi-isomorphism.

Combining [Pri5, Proposition 3.12] with [Pri3, Theorem 4.15 and Corollary 6.35], every strongly quasi-compact derived Artin N -stack over R can be resolved by a derived DM hypergroupoid (a form of homotopy formally étale cosimplicial diagram) in $DG^+ dg\mathrm{CAlg}(R)$.

1.2. Polyvectors. We now fix a chain CDGA R over \mathbb{Q} .

Assumption 1.7. As in [Pri5, §3.3], we now assume that $A \in DG^+dg\mathcal{CAlg}(R)$ has the following properties:

- (1) for any cofibrant replacement $\tilde{A} \rightarrow A$ in the model structure of Lemma 1.3, the morphism $\Omega_{\tilde{A}/R}^1 \rightarrow \Omega_{A/R}^1$ is a levelwise quasi-isomorphism,
- (2) the $A^\#$ -module $(\Omega_{A/R}^1)^\#$ in graded chain complexes is cofibrant (i.e. it has the left lifting property with respect to all surjections of $A^\#$ -modules in graded chain complexes),
- (3) there exists N for which the chain complexes $(\Omega_{A/R}^1 \otimes_A A^0)^i$ are acyclic for all $i > N$.

Of particular interest for us is that these conditions are satisfied when $A = D^*O(X)$ for derived Artin N -hypercgroupoids X . The following is adapted from [Pri5, Definition 3.19] along the lines of [Pri4, Definition 1.3], with the introduction of a dummy variable \hbar of cohomological degree 0.

Definition 1.8. Define the complex of 0-shifted polyvector fields (or strictly speaking, multiderivations) on A by

$$\widehat{\text{Pol}}(A/R, 0) := \prod_{p \geq 0} \widehat{\text{Hom}}_A(\Omega_{A/R}^p, A) \hbar^{p-1}[-p].$$

with graded-commutative multiplication $(a, b) \mapsto \hbar ab$ following the usual conventions for symmetric powers.

The Lie bracket on $\widehat{\text{Hom}}_A(\Omega_{A/R}^1, A)$ then extends to give a bracket (the Schouten–Nijenhuis bracket)

$$[-, -]: \widehat{\text{Pol}}(A/R, 0) \times \widehat{\text{Pol}}(A/R, 0) \rightarrow \widehat{\text{Pol}}(A/R, 0)[-1],$$

determined by the property that it is a bi-derivation with respect to the multiplication operation.

Thus $\widehat{\text{Pol}}(A/R, 0)$ has the natural structure of a P_2 -algebra, and in particular $\widehat{\text{Pol}}(A/R, 0)[1]$ is a differential graded Lie algebra (DGLA) over R .

Definition 1.9. Define a decreasing filtration F on $\widehat{\text{Pol}}(A/R, 0)$ by

$$F^i \widehat{\text{Pol}}(A/R, 0) := \prod_{j \geq i} \widehat{\text{Hom}}_A(\Omega_{A/R}^j, A) \hbar^{j-1}[-j];$$

this has the properties that $\widehat{\text{Pol}}(A/R, 0) = \varprojlim_i \widehat{\text{Pol}}(A/R, 0)/F^i$, with $[F^i, F^j] \subset F^{i+j-1}$, $(\partial \pm \delta)F^i \subset F^i$, and $F^i F^j \subset \hbar^{-1} F^{i+j}$.

Observe that this filtration makes $F^2 \widehat{\text{Pol}}(A/R, n)[1]$ into a pro-nilpotent DGLA.

Definition 1.10. Define the tangent space of polyvectors by

$$T\widehat{\text{Pol}}(A/R, 0) := \widehat{\text{Pol}}(A/R, 0) \oplus \widehat{\text{Pol}}(A/R, 0)\hbar\epsilon,$$

for ϵ of degree 0 with $\epsilon^2 = 0$. Then $T\widehat{\text{Pol}}(A/R, 0)[1]$ is a DGLA with Lie bracket given by $[u + v\epsilon, x + y\epsilon] = [u, x] + [u, y]\epsilon + [v, x]\epsilon$.

Definition 1.11. Given a Maurer–Cartan element $\pi \in \text{MC}(F^2\widehat{\text{Pol}}(A/R, 0)[1])$, define

$$T_\pi\widehat{\text{Pol}}(A/R, 0) := \prod_{p \geq 0} \underline{\widehat{\text{Hom}}}_A(\Omega_{A/R}^p, A)\hbar^p[-p],$$

with derivation $(\partial \pm \delta) + [\pi, -]$ (necessarily square-zero by the Maurer–Cartan conditions).

The product on polyvectors makes this a CDGA (with no need to rescale the product by \hbar), and it inherits the filtration F from $\widehat{\text{Pol}}$.

Given $\pi \in \text{MC}(F^2\widehat{\text{Pol}}(A/R, 0)[1]/F^p)$, we define $T_\pi\widehat{\text{Pol}}(A/R, 0)/F^p$ similarly. This is a CDGA because $F^i \cdot F^j \subset F^{i+j}$.

Regarding $T_\pi\widehat{\text{Pol}}(A/R, 0)[1]$ as an abelian DGLA, observe that $\text{MC}(T_\pi\widehat{\text{Pol}}(A/R, 0)[1])$ is just the fibre of $\text{MC}(T\widehat{\text{Pol}}(A/R, 0)[1]) \rightarrow \text{MC}(\widehat{\text{Pol}}(A/R, 0)[1])$ over π .

1.3. The Hochschild complex of a bi-CDGA.

Definition 1.12. For an A -module M in chain cochain complexes, we define the cohomological Hochschild complex $\text{CC}_R^\bullet(A, M)$ over R as we would for dg algebras, but using the Hom-complexes $\underline{\widehat{\text{Hom}}}$. Thus $\text{CC}_R^\bullet(A, M)$ is the product total complex of a double complex $\underline{\text{CC}}_R^\bullet(A, M)$ given by

$$\underline{\text{CC}}_R^n(A, M) = \underline{\widehat{\text{Hom}}}_R(A^{\otimes R^n}, M),$$

with Hochschild differential $b: \underline{\text{CC}}^{n-1} \rightarrow \underline{\text{CC}}^n$ given by

$$\begin{aligned} (bf)(a_1, \dots, a_n) &= a_1 f(a_2, \dots, a_n) \\ &+ \sum_{i=1}^{n-1} (-1)^i f(a_1, \dots, a_{i-1}, a_i a_{i+1}, a_{i+2}, \dots, a_n) \\ &+ (-1)^n f(a_1, \dots, a_{n-1}) a_n. \end{aligned}$$

There is also a quasi-isomorphic normalised version $N_c \underline{\text{CC}}_R^\bullet(A, M)$, given by the subspaces of functions f with $f(a_1, \dots, a_{i-1}, 1, a_i, \dots, a_n) = 0$ for all i .

We define increasing filtrations F on $\underline{\text{CC}}_R^\bullet(A, M)$ and $\text{CC}_R^\bullet(A, M)$ by good truncation in the Hochschild direction, so $F_p \text{CC}_R^\bullet(A, M) \subset \text{CC}_R^\bullet(A, M)$ is the subspace

$$\prod_{i=0}^{p-1} \underline{\text{CC}}_R^i(A, M)[-i] \times \ker(b: \underline{\text{CC}}_R^p(A, M) \rightarrow \underline{\text{CC}}_R^{p+1}(A, M))[-p].$$

We simply write $\text{CC}_R^\bullet(A)$ and $\underline{\text{CC}}_R^\bullet(A)$ for $\text{CC}_R^\bullet(A, A)$ and $\underline{\text{CC}}_R^\bullet(A, A)$.

Lemma 1.13. *There is a natural brace algebra structure on $\text{CC}_R^\bullet(A)$ over R , compatible with the filtration F . In particular, $\text{CC}_R^\bullet(A)[1]$ is a filtered DGLA over R . On the associated graded brace algebra $\text{gr}^F \text{CC}_R^\bullet(A)$, the Lie bracket and higher braces vanish, and there is a surjective quasi-isomorphism*

$$\text{gr}^F \text{CC}_R^\bullet(A) \rightarrow \widehat{\text{Tot}}(\text{HH}_{R\#}^*(A_{\#}^{\#}), \partial \pm \delta)$$

of brace algebras, where we set all the braces to be 0 on HH^* .

Proof. As in [Vor, §3], there is a brace algebra structure on $\text{CC}_R^\bullet(A)$, with cup product \cdot of cohomological degree 0 and brace operations $(f, g_1, \dots, g_n) \mapsto \{f\}\{g_1, \dots, g_n\}$ of degree cohomological $-n$. Writing $[f, g] := \{f\}\{g\} - (-1)^{\deg f \deg g} \{g\}\{f\}$ defines a Lie bracket of degree -1 , making $\text{CC}_R^n(A)[1]$ into a DGLA.

Compatibility of b with the bracket then implies that $[F_p, F_q] \subset F_{p+q-1}$, and degree considerations also give

$$F_p \cdot F_q \subset F_{p+q}, \quad \{F_p\}\{F_{r_1}, \dots, F_{r_n}\} \subset F_{p+r+1-n},$$

where $r = \sum r_i$. Since $F_{p+r+1-n} \subset F_{p+r}$, this ensures that $(\mathrm{CC}_R^\bullet(A), F)$ is a filtered brace algebra; the bracket vanishes on gr^F , as do the braces for $n \geq 2$.

Since F is defined as good truncation in the Hochschild direction, Hochschild cohomology HH^* is automatically a quasi-isomorphic quotient of gr^F . Any operation of negative degree necessarily vanishes on this quotient, so the quotient map is a brace algebra morphism. \square

Lemma 1.14. *There is an involutive map $i: \mathrm{CC}_R^\bullet(A)[1] \rightarrow \mathrm{CC}_R^\bullet(A)[1]$ of DGLAs given by*

$$i(f)(a_1, \dots, a_m) = -(-1)^{\sum_{i < j} \deg a_i \deg a_j} (-1)^{m(m+1)/2} f(a_m, \dots, a_1).$$

This involution corresponds under the HKR isomorphism to the involution of $\widehat{\mathrm{Pol}}(A, 0)$ which acts on $\underline{\mathrm{Hom}}_A(\Omega_{A/R}^p, A)$ as scalar multiplication by $(-1)^{p-1}$.

Proof. The first statement is proved in [Bra, §2.1], taking the trivial involution on A . For the second statement, given $\phi \in \underline{\mathrm{Hom}}_A(\Omega_{A/R}^p, A)$, the corresponding element f of $\mathrm{CC}^p(A)$ is given by $f(a_1, \dots, a_p) := \phi(da_1 \wedge \dots \wedge da_p)$, and then

$$\begin{aligned} i(f)(a_1, \dots, a_p) &= -(-1)^{\sum_{i < j} \deg a_i \deg a_j} (-1)^{p(p+1)/2} \phi(da_p \wedge \dots \wedge da_1) \\ &= -(-1)^{p(p+1)/2} (-1)^{p(p-1)/2} \phi(da_1 \wedge \dots \wedge da_p) \\ &= -(-1)^p f(a_1, \dots, a_p). \end{aligned}$$

\square

1.4. Quantised 0-shifted polyvectors and quantisations.

Definition 1.15. Define the complex of quantised 0-shifted polyvector fields on A by

$$Q\widehat{\mathrm{Pol}}(A, 0) := \prod_{p \geq 0} F_p \mathrm{CC}_R^\bullet(A) \hbar^{p-1}.$$

Properties of the filtration F from Lemmas 1.13 ensure that $Q\widehat{\mathrm{Pol}}(A, 0)[1]$ is a DGLA.

Definition 1.16. Define a decreasing filtration \tilde{F} on $Q\widehat{\mathrm{Pol}}(A, 0)$ by the subcomplexes

$$\tilde{F}^i Q\widehat{\mathrm{Pol}}(A, 0) := \prod_{j \geq i} F_j \mathrm{CC}_R^\bullet(A) \hbar^{j-1}.$$

This filtration is complete and Hausdorff, with $[\tilde{F}^i, \tilde{F}^j] \subset \tilde{F}^{i+j-1}$. In particular, this makes $\tilde{F}^2 Q\widehat{\mathrm{Pol}}(A, 0)[1]$ into a pro-nilpotent filtered DGLA.

Definition 1.17. Define an E_1 quantisation of A over R to be a Maurer–Cartan element

$$\Delta \in \mathrm{MC}(\tilde{F}^2 Q\widehat{\mathrm{Pol}}(A, 0)).$$

When $A = A_\bullet^0$ is just a CDGA, this gives a curved A_∞ -algebra structure A' on $A[[\hbar]]$ with $A'/\hbar = A$, because $\hbar \mid \Delta$. For more general bi-CDGAs, the stacky and derived structures interact in a non-trivial way on quantisations, and indeed for Poisson structures.

Remark 1.18. To strengthen the analogy between this construction and [Pri4], we could replace $N_c \text{CC}^\bullet(A)$ with its quasi-isomorphic subcomplex of polydifferential operators. The filtration F is then quasi-isomorphic to the order filtration for polydifferential operators, but the latter does not interact so well with the Lie bracket.

If we wished to consider uncurved A_∞ -algebra deformations without inner automorphisms, we would have to replace $\text{CC}^\bullet(A)$ with its sub-DGLA $\ker(\text{CC}_R^\bullet(A) \rightarrow \hat{\text{Tot}} A)$. The analogue for [Pri4] is the kernel of the map $\mathcal{D}_A \rightarrow A$ given by evaluating at 1. As in [Pri4, Remark 1.13], this means that the E_0 analogue of a strict quantisation is a BV algebra deformation.

Definition 1.19. Given a DGLA L , define the Maurer–Cartan set by

$$\text{MC}(L) := \{\omega \in L^1 \mid d\omega + \frac{1}{2}[\omega, \omega] = 0 \in L^2\}.$$

Following [Hin], define the Maurer–Cartan space $\underline{\text{MC}}(L)$ (a simplicial set) of a nilpotent DGLA L by

$$\underline{\text{MC}}(L)_n := \text{MC}(L \otimes_{\mathbb{Q}} \Omega^\bullet(\Delta^n)),$$

where

$$\Omega^\bullet(\Delta^n) = \mathbb{Q}[t_0, t_1, \dots, t_n, \delta t_0, \delta t_1, \dots, \delta t_n] / (\sum t_i - 1, \sum \delta t_i)$$

is the commutative dg algebra of de Rham polynomial forms on the n -simplex, with the t_i of degree 0.

Definition 1.20. We now define another decreasing filtration G on $\widehat{Q\text{Pol}}(A, 0)$ by setting

$$G^i \widehat{Q\text{Pol}}(A, 0) := \hbar^i \widehat{Q\text{Pol}}(A, 0).$$

We then set $G^i \tilde{F}^p := G^i \cap \tilde{F}^p$.

Definition 1.21. Define the space $Q\mathcal{P}(A, 0)$ of E_1 quantisations of A over R to be given by the simplicial set

$$Q\mathcal{P}(A, 0) := \varprojlim_i \underline{\text{MC}}(\tilde{F}^2 \widehat{Q\text{Pol}}(A, 0)[1] / \tilde{F}^{i+2}).$$

Also write

$$Q\mathcal{P}(A, 0) / G^k := \varprojlim_i \underline{\text{MC}}(\tilde{F}^2 \widehat{Q\text{Pol}}(A, 0)[1] / (\tilde{F}^{i+2} + G^k)),$$

so $Q\mathcal{P}(A, 0) = \varprojlim_k Q\mathcal{P}(A, 0) / G^k$.

When R and $A = A^0$ are concentrated in non-negative homological degrees, we can interpret $Q\mathcal{P}(A, 0)$ as a space of deformations of A as an R -linear dg category up to quasi-equivalence, and in general when $A = A^0$ and has bounded cohomology, [LdB] interprets $Q\mathcal{P}(A, 0)$ as a space of deformations of A as an R -linear dg category up to derived Morita equivalence.

For bi-CDGAs, good behaviour of the functor $\hat{\text{Tot}}$ gives us a natural map $\text{CC}_R^\bullet(A) \rightarrow \text{CC}_R^\bullet(\hat{\text{Tot}} A)$ (rarely an equivalence), so E_1 quantisations give rise to curved A_∞ deformations of the CDGA $\hat{\text{Tot}} A$. We now give a stronger statement.

Definition 1.22. If $A \in DG^+ dg\text{CAlg}(R)$, define the bi-dg category $\mathcal{P}er(A)$ as follows. Objects are A -modules M in chain cochain complexes for which $M^\#$ is cofibrant as a graded chain complex over $A^\#$, M^0 is perfect over A^0 , and the map $M^0 \otimes_{A^0} A^\# \rightarrow M^\#$

is a levelwise quasi-isomorphism. Morphisms are given by the chain cochain complexes $\mathcal{H}om_A(M, N)$.

We then define $\text{per}_{dg}(A)$ to have the same objects as $\mathcal{P}er(A)$, and morphisms $\hat{\mathcal{H}om}_A(M, N)$.

For every $M \in \text{per}_{dg}(A)$, we have a $\hat{\text{Tot}} A$ -module $\hat{\text{Tot}} M$, but this need not be cofibrant or perfect. For instance, given $b \in Z_0 Z^1 A$, we may set A_b to be the chain cochain complex $A_\bullet^\#$ with cochain differential $\partial_A + b$. Since $A_b^\# = A^\#$, it lies in $\mathcal{P}er(A)$, but $\hat{\text{Tot}} A_b$ is seldom cofibrant.

Proposition 1.23. *For $A \in DG^+ dg\text{CAlg}(R)$, there is a natural map in the ∞ -category of simplicial sets from $Q\mathcal{P}(A, 0)$ to the space of curved A_∞ deformations $(\text{per}_{dg}(A)[[\hbar]], \{m^{(i)}\}_{i \geq 0})$ of the dg category $\text{per}_{dg}(A)$, with $\hbar^{i-1} \mid m^{(i)}$ for $i \geq 3$.*

Proof. For any R -linear bi-dg category \mathcal{B} , we have a Hochschild complex built from the spaces

$$\underline{\text{CC}}_R^n(\mathcal{B}) = \prod_{x_0, \dots, x_n \in \mathcal{B}} \hat{\mathcal{H}om}_R(\mathcal{B}(x_0, x_1) \otimes_R \dots \otimes_R \mathcal{B}(x_{n-1}, x_n), \mathcal{B}(x_0, x_n)),$$

with $Q\mathcal{P}(\mathcal{B}, 0)$ defined analogously. Properties of $\hat{\text{Tot}}$ then give us a natural map from $Q\mathcal{P}(\mathcal{B}, 0)$ to $Q\mathcal{P}(\hat{\text{Tot}} \mathcal{B}, 0)$, which is the space of curved A_∞ deformations $((\hat{\text{Tot}} \mathcal{B})[[\hbar]], \{m^{(i)}\}_{i \geq 0})$ of the dg category $\hat{\text{Tot}} \mathcal{B}$ with $\hbar^{i-1} \mid m^{(i)}$ for $i \geq 3$; the Maurer–Cartan conditions ensure that $\hbar^i \mid bm^{(i)}$, so every such m does lie in the appropriate piece of the good truncation filtration.

It therefore suffices to show that the map $Q\mathcal{P}(\mathcal{P}er(A), 0) \rightarrow Q\mathcal{P}(A, 0)$ given by restriction to the object $A \in \mathcal{P}er(A)$ is a weak equivalence. By the theory of pro-nilpotent DGLAs, this will follow if $\underline{\text{CC}}_R^n(\mathcal{P}er(A)) \rightarrow \underline{\text{CC}}_R^n(A)$ is a filtered quasi-isomorphism.

We now observe that for any A -linear bi-dg category \mathcal{B} with cofibrant $\mathcal{H}om$ -bicomplexes, there is a spectral sequence

$$\text{HH}_{R\#}^i(A_\#^\#, \text{HH}_{A\#}^j(\mathcal{B}_\#^\#)) \implies \text{HH}_{R\#}^{i+j}(\mathcal{B}_\#^\#).$$

When \mathcal{B} is homotopy Cartesian in the sense that the map $\mathcal{B}^0 \otimes_{A^0} A^\# \rightarrow \mathcal{B}^\#$ is a levelwise quasi-isomorphism, we have a quasi-isomorphism $(\text{HH}_{A\#}^j(\mathcal{B}_\#^0, (\mathcal{B}^0 \otimes_{A^0} A^\#)_\#), \delta) \simeq (\text{HH}_{A\#}^j(\mathcal{B}_\#^\#, \delta)$. We then note that when $\mathcal{B}_\#^0$ is Morita equivalent to $A_\#^0$ as a graded category, the map $M \rightarrow \text{HH}_{A\#}^*(\mathcal{B}_\#^0, (\mathcal{B}_\#^0 \otimes_{A\#}^{\mathbf{L}} M))$ is an isomorphism of graded modules for all $A_\#^0$ -modules M .

Putting these together gives quasi-isomorphisms

$$\text{gr}_j^F \text{CC}_R^\bullet(A) \simeq \text{gr}_j^F \text{CC}_R^\bullet(\mathcal{P}er(A)) :$$

the bi-dg category $\mathcal{P}er(A)$ is homotopy Cartesian because its objects are; since $\mathcal{P}er(A)_\#^0$ is equivalent to the category of graded projective $A_\#^0$ -modules, it is Morita equivalent to $A_\#^0$. Thus $Q\mathcal{P}(\mathcal{P}er(A), 0) \rightarrow Q\mathcal{P}(A, 0)$ is indeed a weak equivalence. \square

Remark 1.24. In [Pri4], we were able to consider E_0 quantisations not just of the structure sheaf \mathcal{O}_X , but also of line bundles, by constructing a \mathbb{G}_m -action on quantised polyvectors.

Similarly, the methods of this paper can be adapted to study E_1 quantisations of any A -linear bi-dg category \mathcal{B} for which the map $\widehat{\text{Tot}} A \rightarrow \text{gr}^F \text{CC}_A^\bullet(\mathcal{B})$ is a quasi-isomorphism — by analogy, line bundles are A -modules for which the map $\widehat{\text{Tot}} A \rightarrow \mathbf{R}\widehat{\text{Hom}}_A(M, M)$ is a quasi-isomorphism. In particular, we can study étale \mathbb{G}_m -gerbes by establishing $B\mathbb{G}_m$ -equivariance. One way to do this is to consider $Q\mathcal{P}(\mathcal{P}er(A), 0)$ as in the proof of Proposition 1.23, since $\mathcal{P}er(A)$ admits an action of the Picard 2-group and hence a $B\mathbb{G}_m$ -action.

The resulting action is necessarily trivial modulo G^1 , so comes from pro-unipotent L_∞ -automorphisms of $Q\widehat{\text{Pol}}(A, 0)$. Since pro-unipotent L_∞ -automorphisms are exponentials of pro-nilpotent L_∞ -derivations, we will in fact have an action of $B\mathbb{G}_m \otimes_{\mathbb{Z}} \mathbb{Q}$, so a notion of quantisation for $(\mathbb{G}_m \otimes_{\mathbb{Z}} \mathbb{Q})$ -gerbes.

1.5. The centre of a quantisation.

Definition 1.25. Define the filtered tangent space to quantised polyvectors by

$$\begin{aligned} TQ\widehat{\text{Pol}}(A, 0) &:= Q\widehat{\text{Pol}}(A, 0) \oplus \prod_{p \geq 0} F_p \text{CC}_R^\bullet(A) \hbar^p \epsilon, \\ \tilde{F}^j TQ\widehat{\text{Pol}}(A, 0) &:= \tilde{F}^j Q\widehat{\text{Pol}}(A, 0) \oplus \prod_{p \geq j} F_p \text{CC}_R^\bullet(A) \hbar^p \epsilon, \end{aligned}$$

for ϵ of degree 0 with $\epsilon^2 = 0$. Then $TQ\widehat{\text{Pol}}(A, 0)[1]$ is a DGLA, with Lie bracket given by $[u + v\epsilon, x + y\epsilon] = [u, x] + [u, y]\epsilon + [v, x]\epsilon$.

Definition 1.26. Given a Maurer–Cartan element $\Delta \in \text{MC}(\tilde{F}^2 Q\widehat{\text{Pol}}(A, 0)[1])$, define the centre of (A, Δ) by

$$T_\Delta Q\widehat{\text{Pol}}(A, 0) := \prod_{p \geq 0} F_p \text{CC}_R^\bullet(A) \hbar^p,$$

with derivation $\partial \pm \delta \pm b + [\Delta, -]$ (necessarily square-zero by the Maurer–Cartan conditions).

This has a filtration

$$\tilde{F}^i T_\Delta Q\widehat{\text{Pol}}(A, 0) := \prod_{p \geq i} F_p \text{CC}_R^\bullet(A) \hbar^p,$$

making $T_\Delta Q\widehat{\text{Pol}}(A, 0)$ a filtered brace algebra by Lemma 1.13. Given $\Delta \in \text{MC}(F^2 Q\widehat{\text{Pol}}(A, 0)/\tilde{F}^p)$, we define $T_\Delta Q\widehat{\text{Pol}}(A, 0)/\tilde{F}^p$ similarly — this is also a brace algebra as \tilde{F}^p is a brace ideal.

Observe that $T_\Delta Q\mathcal{P}(A, 0) := \underline{\text{MC}}(\tilde{F}^2 T_\Delta Q\widehat{\text{Pol}}(A, 0)[1])$ is just the fibre of $\underline{\text{MC}}(\tilde{F}^2 TQ\widehat{\text{Pol}}(A, 0)[1]) \rightarrow \underline{\text{MC}}(\tilde{F}^2 Q\widehat{\text{Pol}}(A, 0)[1])$ over Δ .

Similarly to Definition 1.20, there are filtrations G on $TQ\widehat{\text{Pol}}(A, 0), T_\Delta Q\widehat{\text{Pol}}(A, 0)$ given by powers of \hbar . Since $\text{gr}_G^i \tilde{F}^{p-i} Q\widehat{\text{Pol}} = \prod_{j \geq p-i} \text{gr}_{j-i}^F \text{CC}_R^\bullet(A) \hbar^{j-1}$, the HKR isomorphism gives maps

$$\begin{aligned} \text{gr}_G^i \tilde{F}^p Q\widehat{\text{Pol}}(A, 0) &\rightarrow \prod_{j \geq p} \widehat{\text{Hom}}_A(\Omega_{A/R}^{j-i}, A) \hbar^{j-1} [i - j] \\ \text{gr}_G^i \tilde{F}^p T_\Delta Q\widehat{\text{Pol}}(A, 0) &\rightarrow \prod_{j \geq p} \widehat{\text{Hom}}_A(\Omega_{A/R}^{j-i}, A) \hbar^j [i - j], \end{aligned}$$

which are quasi-isomorphisms by our hypotheses on A (see Assumption 1.7).

For the filtration F of Definition 1.9, we may rewrite these maps as

$$\begin{aligned} \mathrm{gr}_G^i \tilde{F}^p Q\widehat{\mathrm{Pol}}(A, 0) &\rightarrow F^{p-i} \widehat{\mathrm{Pol}}(A, 0) \hbar^i, \\ \mathrm{gr}_G^i \tilde{F}^p T_\Delta Q\widehat{\mathrm{Pol}}(A, 0) &\rightarrow F^{p-i} T_{\pi_\Delta} \widehat{\mathrm{Pol}}(A, 0) \hbar^i, \end{aligned}$$

where $\pi_\Delta \in \mathrm{MC}(F^2 \widehat{\mathrm{Pol}}(A, 0)[1])$ denotes the image of Δ under the map $\mathrm{gr}_G^0 \tilde{F}^2 Q\widehat{\mathrm{Pol}}(A, 0) \rightarrow F^2 \widehat{\mathrm{Pol}}(A, 0)$.

Since the cohomology groups of $T_{\pi_\Delta} \widehat{\mathrm{Pol}}(A, 0)$ are Poisson cohomology, we will refer to the cohomology groups of $T_\Delta Q\widehat{\mathrm{Pol}}(A, 0)$ as quantised Poisson cohomology.

Definition 1.27. Say that an E_1 quantisation $\Delta = \sum_{j \geq 2} \Delta_j \hbar^{j-1}$ is non-degenerate if the map

$$\Delta_2^\# : \mathrm{Tot}^\Pi(\Omega_{A/R}^1 \otimes_A A^0) \rightarrow \underline{\mathrm{Hom}}_A(\Omega_A^1, A^0)$$

is a quasi-isomorphism and $\mathrm{Tot}^\Pi(\Omega_{A/R}^1 \otimes_A A^0)$ is a perfect complex over A^0 .

Definition 1.28. Define the tangent spaces

$$TQ\mathcal{P}(A, 0) := \varprojlim_i \underline{\mathrm{MC}}(\tilde{F}^2 TQ\widehat{\mathrm{Pol}}(A, 0)[1]/\tilde{F}^{i+2}),$$

with $TQ\mathcal{P}(A, 0)/G^k$, defined similarly.

These are simplicial sets over $Q\mathcal{P}(A, 0)$ (resp. $Q\mathcal{P}(A, 0)/G^k$), fibred in simplicial abelian groups.

Definition 1.29. Define the canonical tangent vector

$$\sigma = -\partial_{\hbar^{-1}} : Q\widehat{\mathrm{Pol}}(A, 0) \rightarrow TQ\widehat{\mathrm{Pol}}(A, 0)$$

by $\alpha \mapsto \alpha + \epsilon \hbar^2 \frac{\partial \alpha}{\partial \hbar}$. Note that this is a morphism of filtered DGLAs, so gives a map $\sigma : Q\mathcal{P}(A, 0) \rightarrow TQ\mathcal{P}(A, 0)$, with $\sigma(\Delta) \in Z^2(\tilde{F}^2 T_\Delta Q\widehat{\mathrm{Pol}}(A, 0))$.

1.6. Self-dual quantisations.

Definition 1.30. Define an involution $Q\widehat{\mathrm{Pol}}(A, 0) \xrightarrow{*} Q\widehat{\mathrm{Pol}}(A, 0)$ by $\Delta^*(\hbar) := i(\Delta)(-\hbar)$, for the involution i of Lemma 1.14.

Definition 1.31. Lemma 1.14 ensures that $*$ is a morphism of DGLAs, and we define the space $Q\mathcal{P}(A, 0)^{sd} \subset Q\mathcal{P}(A, 0)$ of self-dual quantisations to be the fixed points of the involution $*$. This inherits cofiltrations \tilde{F} and G from $Q\mathcal{P}(A, 0)$.

In particular, this means that when $A = A^0$ is concentrated in degree zero, elements of $Q\mathcal{P}(A, 0)^{sd}$ can be represented by associative algebra deformations (A', \star_\hbar) of A , with

$$a \star_{-\hbar} b = b \star_\hbar a.$$

More generally, when R and $A = A^0$ are concentrated in non-negative homological degrees, elements of $Q\mathcal{P}(A, 0)^{sd}$ are algebroid quantisations of A equipped with a contravariant involution which is semilinear under the transformation $\hbar \mapsto -\hbar$.

For general bi-CDGAs A , $Q\mathcal{P}(A, 0)^{sd}$ gives rise via Proposition 1.23 to curved A_∞ -deformations $\tilde{\mathcal{P}}_\hbar$ of the dg category $\mathrm{per}_{\mathrm{dg}}(A)$ of perfect complexes, equipped with an involution $\tilde{\mathcal{P}}_{-\hbar} \simeq \tilde{\mathcal{P}}_\hbar^{\mathrm{opp}}$ lifting the duality functor $\mathcal{H}om_A(-, A)$ on $\mathrm{per}_{\mathrm{dg}}(A)$.

Remark 1.32. As in Remark 1.24, we may also consider self-duality for \mathbb{G}_m -gerbes. Since the functor sending a gerbe to its opposite is just given by the inversion map on $B^2\mathbb{G}_m$, involutive gerbes are classified by $B^2\mu_2$, the homotopy fixed points of the inversion map.

However, as observed in Remark 1.24, the space of quantisations over $B^2\mathbb{G}_m$ is the pullback of a space over $B^2(\mathbb{G}_m \otimes_{\mathbb{Z}} \mathbb{Q})$, so the space of self-dual quantisations over $B^2\mu_2$ is constant. This means that to every self-dual quantisation of A there correspond self-dual quantisations of all μ_2 -gerbes, and in particular of $\text{per}_{\text{dg}}(A)$ with duality functor $\mathbf{R}\mathcal{H}om(-, \mathcal{L})$ for any line bundle \mathcal{L} . One way to make sense of this example is that even if \mathcal{L} does not have a square root, there is necessarily an automorphism of the Hochschild complex acting as a square root of \mathcal{L} , and thus intertwining between the respective duality functors.

Lemma 1.33. *There are canonical weak equivalences*

$$\begin{aligned} Q\mathcal{P}(A, 0)^{sd}/G^{2i} &\rightarrow Q\mathcal{P}(A, 0)^{sd}/G^{2i-1} \\ Q\mathcal{P}(A, 0)^{sd}/G^{2i+1} &\rightarrow (Q\mathcal{P}(A, 0)^{sd}/G^{2i}) \times_{(Q\mathcal{P}(A, 0)/G^{2i})}^h (Q\mathcal{P}(A, 0)/G^{2i+1}). \end{aligned}$$

Proof. This follows in much the same way as [Pri4, Lemma 4.5]. Lemma 1.14 ensures that the involution $*$ acts trivially on $\widehat{\text{Pol}}(A, 0)$, since it maps $f\hbar^{p-1}$ to $(-1)^{p-1}(-\hbar)^{p-1}$ for $f \in \widehat{\text{Hom}}_A(\Omega_{A/R}^p, A)\hbar^{p-1}$. It therefore acts as multiplication by $(-1)^k$ on $\text{gr}_G^k \widehat{\text{Pol}}(A, 0) = \hbar^k \text{gr}_G^0 \widehat{\text{Pol}}(A, 0)$, giving quasi-isomorphisms

$$\text{gr}_G^k \tilde{F}^p Q\widehat{\text{Pol}}(A, 0)^{sd} \simeq \begin{cases} \text{gr}_G^k \tilde{F}^p Q\widehat{\text{Pol}}(A, 0) & k \text{ even} \\ 0 & k \text{ odd.} \end{cases}$$

The results then follow from the fibre sequences

$$Q\mathcal{P}(A, 0)^{sd}/G^{k+1} \rightarrow Q\mathcal{P}(A, 0)^{sd}/G^k \rightarrow \underline{\text{MC}}(\text{gr}_G^k \tilde{F}^2 Q\widehat{\text{Pol}}(A, 0)^{sd}[2])$$

coming from obstruction theory for abelian extensions of DGLAs. \square

In particular, Lemma 1.33 gives $Q\mathcal{P}(A, 0)^{sd}/G^2 \simeq Q\mathcal{P}(A, 0)^{sd}/G^1 \simeq \mathcal{P}(A, 0)$, so every unshifted Poisson structure admits an essentially unique first-order self-dual quantisation.

2. QUANTISATIONS AND DE RHAM POWER SERIES

Recall that we are fixing a chain CDGA R over \mathbb{Q} , and a cofibrant bi-CDGA A over R . We denote the chain differentials on A and R by δ , and the cochain differential on A by ∂ .

2.1. Generalised pre-symplectic structures. We now adapt some definitions from [Pri4, §1.2] and [Pri5, §3.3.2].

Definition 2.1. Define the de Rham complex $\text{DR}(A/R)$ to be the product total complex of the bicomplex

$$\text{Tot}^{\Pi} A \xrightarrow{d} \text{Tot}^{\Pi} \Omega_{A/R}^1 \xrightarrow{d} \text{Tot}^{\Pi} \Omega_{A/R}^2 \xrightarrow{d} \dots,$$

so the total differential is $d \pm \partial \pm \delta$.

We define the Hodge filtration F on $\text{DR}(A/R)$ by setting $F^p \text{DR}(A/R) \subset \text{DR}(A/R)$ to consist of terms $\text{Tot}^{\Pi} \Omega_{A/R}^i$ with $i \geq p$. In particular, $F^p \text{DR}(A/R) = \text{DR}(A/R)$ for $p \leq 0$.

Definition 2.2. When A is a cofibrant bi-CDGA over R , recall that a 0-shifted pre-symplectic structure ω on A/R is an element

$$\omega \in Z^2 F^2 \mathrm{DR}(A/R).$$

It is called symplectic if $\omega_2 \in Z^2 \mathrm{Tot}^{\Pi} \Omega_{A/R}^2$ induces a quasi-isomorphism

$$\omega_2^{\sharp}: \underline{\mathrm{Hom}}_A(\Omega_{A/R}^1, A^0) \rightarrow \mathrm{Tot}^{\Pi}(\Omega_{A/R}^1 \otimes_A A^0)$$

and $\mathrm{Tot}^{\Pi}(\Omega_{A/R}^1 \otimes_A A^0)$ is a perfect complex over A^0 .

Definition 2.3. Define a decreasing filtration \tilde{F} on $\mathrm{DR}(A/R)[[\hbar]]$ by

$$\tilde{F}^p \mathrm{DR}(A/R) := \prod_{i \geq 0} F^{p-i} \mathrm{DR}(A/R) \hbar^i.$$

Define a further filtration G by $G^k \mathrm{DR}(A/R)[[\hbar]] = \hbar^k \mathrm{DR}(A/R)[[\hbar]]$.

Definition 2.4. Define the space of generalised 0-shifted pre-symplectic structures on A/R to be the simplicial set

$$\mathrm{GPreSp}(A/R, 0) := \varprojlim_i \underline{\mathrm{MC}}(\tilde{F}^2 \mathrm{DR}(A/R)[[\hbar]][1]/\tilde{F}^{i+2}),$$

where we regard the cochain complex $\mathrm{DR}(A/R)[1]$ as a DGLA with trivial bracket. Write $\mathrm{PreSp} = \mathrm{GPreSp}/G^1$.

Also write $\mathrm{GPreSp}(A/R, 0)/\hbar^k := \varprojlim_i \underline{\mathrm{MC}}((\tilde{F}^2 \mathrm{DR}(A/R)[\hbar][1]/(G^k + \tilde{F}^{i+2})),$ so $\mathrm{GPreSp}(A/R, 0) = \varprojlim_k \mathrm{GPreSp}(A/R, 0)/\hbar^k$.

Set $G\mathrm{Sp}(A/R, 0) \subset \mathrm{GPreSp}(A/R, 0)$ to consist of the points whose images in $\mathrm{PreSp}(A/R, 0)$ are symplectic structures — this is a union of path-components.

Remarks 2.5. Note that Definition 2.4 is not the obvious analogue of the definition of generalised (-1) -shifted pre-symplectic structures from [Pri4, Definition 1.29], which used the convolution $(G * \tilde{F})^2 = \tilde{F}^2 + G^1$ in place of \tilde{F}^2 for reasons specific to negatively shifted structures. The only difference lies in the linear term, which is where the correspondence between generalised symplectic structures and non-degenerate quantisations breaks down anyway — replacing \tilde{F}^2 with $(G * \tilde{F})^2$ would not significantly affect the main results of either paper, nor would eliminating the linear term altogether.

Also note that $\mathrm{GPreSp}(A/R, 0)$ is canonically weakly equivalent to the Dold–Kan denormalisation of the good truncation complex $\tau^{\leq 0}(\tilde{F}^2 \mathrm{DR}(A/R)[[\hbar]][2])$ (and similarly for the various quotients we consider), but the description in terms of $\underline{\mathrm{MC}}$ will simplify comparisons. In particular, we have

$$\pi_i \mathrm{GPreSp}(A/R, 0) \cong H^{2-i}(F^2 \mathrm{DR}(A/R)) \times \hbar H^{2-i}(F^1 \mathrm{DR}(A/R)) \times \hbar^2 H^{2-i}(\mathrm{DR}(A/R))[[\hbar]].$$

2.2. Formality.

Definition 2.6. Write GT for the Grothendieck–Teichmüller group. This is an affine group scheme over \mathbb{Q} , with reductive quotient \mathbb{G}_m . Denote the pro-unipotent radical $\ker(\mathrm{GT} \rightarrow \mathbb{G}_m)$ by GT^1 .

Write $\mathrm{Levi}_{\mathrm{GT}}$ for the space of Levi decompositions of GT , i.e. sections of $\mathrm{GT} \rightarrow \mathbb{G}_m$. By the general theory of pro-algebraic groups in characteristic 0, the space $\mathrm{Levi}_{\mathrm{GT}}$ is an affine scheme over \mathbb{Q} equipped with the structure of a trivial GT^1 -torsor via the adjoint action.

For any $\lambda \in k^\times$, a Levi decomposition $w \in \text{Levi}_{\text{GT}}(k)$ gives a λ -Drinfeld associator $w(\lambda) \in \text{GT}(k)$. When λ is not a root of unity, this map from $\text{Levi}_{\text{GT}}(k)$ to the set of λ -Drinfeld associators is an isomorphism, since both sets are $\text{GT}^1(k)$ -torsors.

As explained succinctly in [Pet], formality of the \mathbb{Q} -linear E_2 operad is a consequence of the observation that the Grothendieck–Teichmüller group is a pro-unipotent extension of \mathbb{G}_m . Since GT acts on E_2 , any Levi decomposition $w: \mathbb{G}_m \rightarrow \text{GT}$ gives a weight decomposition (i.e. a \mathbb{G}_m -action) of E_2 which splits the good truncation filtration, so gives an equivalence between E_2 and P_2 respecting the natural map from the Lie operad.

Definition 2.7. Given a Levi decomposition $w \in \text{Levi}_{\text{GT}}(\mathbb{Q})$, we denote by p_w the resulting ∞ -functor from E_2 -algebras to P_2 -algebras over \mathbb{Q} , which respects the underlying L_∞ -algebras.

As in [Vor], brace algebras are naturally E_2 -algebras, so $\text{CC}_R^\bullet(A)$ has an E_2 -algebra structure. Moreover, the equivalence between E_2 and P_2 necessarily respects the good truncation filtrations, and the filtered complex $(\text{CC}_R^\bullet(A), F)$ is an algebra with respect to the brace operad filtered by good truncation. This yields a filtered P_2 -algebra $(p_w \text{CC}_R^\bullet(A), F)$ over A with $F_p \cdot F_q \subset F_{p+q}$ and $[F_p, F_q] \subset F_{p+q-1}$, with a filtered L_∞ -quasi-isomorphism $(p_w \text{CC}_R^\bullet(A)[1], F) \simeq (\text{CC}_R^\bullet(A)[1], F)$.

Definition 2.8. For any of the definitions from §1, we add the subscript w to indicate that we are replacing $(\text{CC}_R^\bullet(A), F)$ with $(p_w \text{CC}_R^\bullet(A), F)$ in the construction.

Since these DGLAs are quasi-isomorphic and $\underline{\text{MC}}$ preserves weak equivalences, in particular we have canonical weak equivalences $Q\mathcal{P}_w(A, 0) \simeq Q\mathcal{P}(A, 0)$. Properties of the filtration \tilde{F} then ensure that the complexes $T_\Delta Q\widehat{\text{Pol}}_w(A, 0)$ are filtered P_2 -algebras.

Remark 2.9. Rather than just choosing $w \in \text{Levi}_{\text{GT}}(\mathbb{Q})$, a more natural approach might be to consider the simplicial set $\mathbf{RLevi}_{\text{GT}}(R)$ of all Levi decompositions over R . This would lead to a space $Q\mathcal{P}_{\text{Levi}}(A, 0)$ over $\mathbf{RLevi}_{\text{GT}}(R)$ with fibre $Q\mathcal{P}_w(A, 0)$ over w and a canonical weak equivalence $Q\mathcal{P}_{\text{Levi}}(A, 0) \simeq \mathbf{RLevi}_{\text{GT}}(R) \times Q\mathcal{P}(A, 0)$.

2.3. Compatible quantisations. We will now develop the notion of compatibility between a generalised pre-symplectic structure and an E_1 quantisation, generalising the notion of compatibility between 0-shifted pre-symplectic and Poisson structures from [Pri5]. The following definitions are adapted from [Pri5, Definition 1.16].

Definition 2.10. Given a bi-CDGA B over A and a derivation $\Delta \in \text{MC}(\widehat{\text{Hom}}_B(\Omega_{B/R}^1, B))$, define

$$\mu(-, \Delta): \text{DR}(A/R) \rightarrow \widehat{\text{Tot}} B$$

to be the morphism of graded A -algebras given on generators $\Omega_{A/R}^1$ by setting

$$\mu(\text{adf}, \Delta) := a\Delta(f),$$

and then applying $\widehat{\text{Tot}}$ (noting that $\text{Tot}^\Pi \Omega_A^p = \widehat{\text{Tot}} \Omega_A^p$).

The proof of [Pri5, Lemma 1.17] ensures that this becomes a chain map (and hence an R -CDGA morphism)

$$\mu(-, \Delta): \text{DR}(A/R) \rightarrow (\widehat{\text{Tot}} B, (\partial \pm \delta)_B + \Delta).$$

Definition 2.11. Given a choice $w \in \text{Levi}_{\text{GT}}(\mathbb{Q})$ of Levi decomposition for GT and $\Delta \in Q\mathcal{P}(A, 0)_w/G^j$ define

$$\mu_w(-, \Delta): \text{DR}(A/R)[[\hbar]]/\hbar^j \rightarrow T_\Delta Q\widehat{\text{Pol}}_w(A, 0)/G^j$$

by applying Definition 2.10 to the bi-CDGAs

$$T_0 Q\widehat{\text{Pol}}(A, 0) := \left(\prod_{i=0}^k F_i p_w \underline{\text{CC}}_R^\bullet(A) \hbar^i \right) / G^j$$

and the derivation $[\Delta, -]$, then taking the limit over all k . Observe that this map preserves the filtration \tilde{F} .

Definition 2.12. We say that a generalised pre-symplectic structure ω and an E_1 quantisation Δ are w -compatible (or a w -compatible pair) if

$$[\mu_w(\omega, \Delta)] = [-\partial_{\hbar^{-1}}(\Delta)] \in H^1(\tilde{F}^2 T_\Delta Q\widehat{\text{Pol}}_w(A, 0)) \cong H^1(\tilde{F}^2 T_\Delta Q\widehat{\text{Pol}}(A, 0)),$$

where $\sigma = -\partial_{\hbar^{-1}}$ is the canonical tangent vector of Definition 1.29.

Definition 2.13. Given a simplicial set Z , an abelian group object A in simplicial sets over Z , a space X over Z and a morphism $s: X \rightarrow A$ over Z , define the homotopy vanishing locus of s over Z to be the homotopy limit of the diagram

$$X \begin{array}{c} \xrightarrow{s} \\ \rightrightarrows \\ \xrightarrow{0} \end{array} A \longrightarrow Z.$$

Definition 2.14. Define the space $Q\text{Comp}_w(A, 0)$ of w -compatible quantised 0-shifted pairs to be the homotopy vanishing locus of

$$(\mu_w - \sigma): G\text{PreSp}(A/R, 0) \times Q\mathcal{P}_w(A, 0) \rightarrow TQ\mathcal{P}_w(A, 0)$$

over $Q\mathcal{P}_w(A, 0)$

We define a cofiltration on this space by setting $Q\text{Comp}_w(A, 0)/G^j$ to be the homotopy vanishing locus of

$$(\mu_w - \sigma): (G\text{PreSp}(A/R, 0)/G^j) \times (Q\mathcal{P}_w(A, 0)/G^j) \rightarrow TQ\mathcal{P}_w(A, 0)/G^j$$

over $Q\mathcal{P}_w(A, 0)/G^j$.

When $j = 1$, note that this recovers the notion of compatible 0-shifted pairs from [Pri5, §3.3.3].

Definition 2.15. Define $Q\text{Comp}_w(A, 0)^{\text{nondeg}} \subset Q\text{Comp}_w(A, 0)$ to consist of w -compatible quantised pairs (ω, Δ) with Δ non-degenerate. This is a union of path-components, and by [Pri5, Lemma 1.22] any pre-symplectic form compatible with a non-degenerate quantisation is symplectic, so there is a natural projection

$$Q\text{Comp}_w(A, 0)^{\text{nondeg}} \rightarrow G\text{Sp}(A/R, 0)$$

as well as the canonical map

$$Q\text{Comp}_w(A, 0)^{\text{nondeg}} \rightarrow Q\mathcal{P}_w(A, 0)^{\text{nondeg}}.$$

2.4. The equivalences.

Proposition 2.16. *For any Levi decomposition w of GT, the canonical map*

$$Q\text{Comp}_w(A, 0)^{\text{nondeg}} \rightarrow Q\mathcal{P}_w(A, 0)^{\text{nondeg}} \simeq Q\mathcal{P}(A, 0)^{\text{nondeg}}$$

is a weak equivalence. In particular, there is a morphism

$$Q\mathcal{P}(A, 0)^{\text{nondeg}} \rightarrow G\text{Sp}(A/R, 0)$$

in the homotopy category of simplicial sets.

Proof. We adapt the proof of [Pri5, Proposition 1.26]. For any $\Delta \in Q\mathcal{P}_w(A, 0)$, the homotopy fibre of $Q\text{Comp}_w(A/R, 0)^{\text{nondeg}}$ over Δ is just the homotopy fibre of

$$\mu_w(-, \Delta): G\text{PreSp}(A/R, 0) \rightarrow T_\Delta Q\mathcal{P}_w(A, 0)$$

over $-\partial_{\hbar^{-1}}(\Delta)$.

The map $\mu_w(-, \Delta): \text{DR}(A/R)[[\hbar]] \rightarrow T_\Delta \widehat{Q\text{Pol}}_w(A, 0)$ is a morphism of complete \tilde{F} -filtered $R[[\hbar]]$ -CDGAs by the proof of [Pri5, Lemma 1.17]. Since the morphism is $R[[\hbar]]$ -linear, it maps $G^k \tilde{F}^p \text{DR}(A/R)[[\hbar]]$ to $G^k \tilde{F}^p T_\Delta \widehat{Q\text{Pol}}_w(A, 0)$. Non-degeneracy of Δ_2 modulo F^1 implies that $\mu_w(-, \Delta)$ induces quasi-isomorphisms

$$\text{Tot}^\Pi \Omega^{p-k} \hbar^k [k-p] \rightarrow \widehat{\text{Hom}}_A(\Omega_{A/R}^{p-k}, A) \hbar^{p-k} [k-p]$$

on the associated gradeds $\text{gr}_G^k \text{gr}_{\tilde{F}}^p$. We therefore have a quasi-isomorphism of bifiltered complexes, so we have isomorphisms on homotopy groups:

$$\begin{aligned} \pi_j G\text{PreSp}(A/R, 0) &\rightarrow \pi_j T_\Delta Q\mathcal{P}(A, 0) \\ H^{2-j}(\tilde{F}^2 \text{DR}(A/R)[[\hbar]]) &\rightarrow H^{2-j}(\tilde{F}^2 T_\Delta \widehat{Q\text{Pol}}(A, 0)). \end{aligned}$$

□

Proposition 2.17. *For any Levi decomposition w of GT, the maps*

$$\begin{aligned} Q\mathcal{P}_w(A, 0)^{\text{nondeg}}/G^j &\rightarrow (Q\mathcal{P}_w(A, 0)^{\text{nondeg}}/G^2) \times_{(G\text{Sp}(A, 0)/G^2)}^{\hbar} (G\text{Sp}(A, 0)/G^j) \\ &\simeq (Q\mathcal{P}_w(A, 0)^{\text{nondeg}}/G^2) \times \prod_{2 \leq i < j} \underline{\text{MC}}(\text{DR}(A/R) \hbar^i [1]) \end{aligned}$$

coming from Proposition 2.16 are weak equivalences for all $j \geq 2$.

Proof. The proof of [Pri4, Proposition 1.40] generalises to this setting. We have a commutative diagram

$$\begin{array}{ccccc} (Q\text{Comp}_w(A, 0)/G^{j+1})_{(\omega, \pi)} & \longrightarrow & (Q\text{Comp}_w(A, 0)/G^j)_{(\omega, \pi)} & \longrightarrow & \underline{\text{MC}}(N(\omega, \pi, j)[2]) \\ \downarrow & & \downarrow & & \downarrow \\ (G\text{PreSp}(A, 0)/G^{j+1})_\omega & \longrightarrow & (G\text{PreSp}(A, 0)/G^j)_\omega & \longrightarrow & \underline{\text{MC}}(F^{2-j} \hbar^j \text{DR}(A/R)[2]) \end{array}$$

of fibre sequences, with $N(\omega, \pi, j)$ the cocone of the map

$$F^{2-j} \text{DR}(A/R) \hbar^j \oplus (F^{2-j} \widehat{\text{Pol}}(A, 0) \hbar^j, \delta_\pi) \rightarrow F^{2-j} T_\pi \widehat{\text{Pol}}(A, 0) \hbar^j$$

given by combining

$$\mu(-, \pi): F^{2-j} \text{DR}(A/R) \hbar^j \rightarrow F^{2-j} T_\pi \widehat{\text{Pol}}(A, 0) \hbar^j$$

with

$$\nu(\omega, \pi) + \partial_{\hbar^{-1}}: F^{2-j} T_\pi \widehat{\text{Pol}}(A, 0) \hbar^{j-1} \rightarrow F^{2-j} T_\pi \widehat{\text{Pol}}(A, 0) \hbar^j.$$

Here $\nu(\omega, \pi)$ is the tangent map of $\mu(\omega, -)$ at π , given by $\mu(\omega, \pi + \rho\epsilon) = \mu(\omega, \pi) + \nu(\omega, \pi)(\rho)\epsilon$ with $\epsilon^2 = 0$.

As in [Pri4, Lemma 1.39], on the associated graded piece

$$\text{gr}_F^p \widehat{\text{Pol}}(A, 0) \hbar^j = \widehat{\text{Hom}}_A(\Omega_{A/R}^p, A) \hbar^{p+j-1},$$

the map $\nu(\omega, \pi)$ is given by $p\Lambda^p(\pi^\# \circ \omega^\#)\hbar$, while $\partial_{\hbar^{-1}} = (1-j-p)\hbar$. Since π is non-degenerate, $\pi^\# \circ \omega^\#$ is homotopic to 1, so $\text{gr}_F^p(\nu(\omega, \pi) + \partial_{\hbar^{-1}})$ is homotopic to $(1-j)\hbar$. As this is an isomorphism for all $j \geq 2$, the map $N(\omega, \pi, j) \rightarrow F^{2-j} \text{DR}(A/R) \hbar^j$ is

quasi-isomorphism, which inductively gives the required weak equivalences from the fibre sequences above. \square

Remark 2.18. Taking the limit over all j , Proposition 2.17 gives an equivalence

$$Q\mathcal{P}_w(A, 0)^{\text{nondeg}} \simeq (Q\mathcal{P}_w(A, 0)^{\text{nondeg}}/G^2) \times \prod_{i \geq 2} \underline{\text{MC}}(\text{DR}(A/R)\hbar^i[1]);$$

in particular, this means that there is a canonical map

$$(Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^2) \rightarrow Q\mathcal{P}(A, 0)^{\text{nondeg}},$$

dependent on w , corresponding to the distinguished point $0 \in \underline{\text{MC}}(\hbar^2 \text{DR}(A/R)[\hbar])$.

Thus to quantise a non-degenerate 0-shifted Poisson structure $\pi = \sum_{j \geq 2} \pi_j$ (or equivalently, by [Pri5, Corollary 1.38], a 0-shifted symplectic structure), it suffices to lift the power series $\sum_{j \geq 2} \pi_j \hbar^{j-1}$ to a Maurer–Cartan element of $\prod_{j \geq 2} (F_j \text{CC}_R^\bullet(A)/F_{j+2})\hbar^{j-1}$.

Even if π is degenerate, a variant of Proposition 2.17 still holds. Because $\pi^\# \circ \omega^\#$ is homotopy idempotent, the map $\text{gr}_F^p \nu(\omega, \pi)$ has eigenvalues in the interval $[0, p]$, so we just replace $(1 - j)$ with an operator having eigenvalues in the interval $[1 - p - j, 1 - j]$. Since this is still a quasi-isomorphism for $j > 1$, we have

$$Q\text{Comp}_w(A, 0) \simeq (Q\text{Comp}_w(A, 0)/G^2) \times \prod_{i \geq 2} \underline{\text{MC}}(\text{DR}(A/R)\hbar^i).$$

giving a sufficient first-order criterion for degenerate quantisations to exist.

Remark 2.19. As in Remark 2.9, we could consider the space $\mathbf{R}\text{Levi}_{\text{GT}}(R)$ of R -linear Levi decompositions, and the proof of Proposition 2.17 then gives equivalences

$$\begin{aligned} & \mathbf{R}\text{Levi}_{\text{GT}}(R) \times Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^j \\ & \rightarrow \mathbf{R}\text{Levi}_{\text{GT}}(R) \times (Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^2) \times_{(G\text{Sp}(A, 0)/G^2)}^h (G\text{Sp}(A, 0)/G^j) \\ & \simeq \mathbf{R}\text{Levi}_{\text{GT}}(R) \times (Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^2) \times \prod_{2 \leq i < j} \underline{\text{MC}}(\text{DR}(A/R)\hbar^i[1]) \end{aligned}$$

over $\mathbf{R}\text{Levi}_{\text{GT}}(R)$.

2.4.1. Self-duality.

Theorem 2.20. *For any Levi decomposition w of GT, there is a canonical weak equivalence*

$$Q\mathcal{P}(A, 0)^{\text{nondeg}, sd} \simeq \mathcal{P}(A, 0)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2 \text{DR}(A/R)[\hbar^2][1]).$$

In particular, w gives a canonical choice of self-dual quantisation for any non-degenerate 0-shifted Poisson structure on A .

Proof. Lemma 1.33 implies that we have weak equivalences

$$\begin{aligned} Q\mathcal{P}(A, 0)^{sd}/G^{2i} & \rightarrow Q\mathcal{P}(A, 0)^{sd}/G^{2i-1} \\ Q\mathcal{P}(A, 0)^{sd}/G^{2i+1} & \rightarrow (Q\mathcal{P}(A, 0)^{sd}/G^{2i}) \times_{(Q\mathcal{P}(A, 0)/G^{2i})}^h (Q\mathcal{P}(A, 0)/G^{2i+1}). \end{aligned}$$

Combined with Proposition 2.17, the latter gives weak equivalences

$$Q\mathcal{P}(A, 0)^{\text{nondeg}, sd}/G^{2i+1} \rightarrow (Q\mathcal{P}(A, 0)^{\text{nondeg}, sd}/G^{2i}) \times \underline{\text{MC}}(\hbar^{2i} \text{DR}(X/R)[1])$$

for all $i > 0$, so

$$\begin{aligned} Q\mathcal{P}(A, 0)^{\text{nondeg}, sd}/G^{2i+1} & \simeq (Q\mathcal{P}(A, 0)^{\text{nondeg}, sd}/G^{2i}) \times \underline{\text{MC}}(\hbar^{2i} \text{DR}(X/R)[1]) \\ & \simeq Q\mathcal{P}(A, 0)^{\text{nondeg}}/G^{2i-1} \times \underline{\text{MC}}(\hbar^{2i} \text{DR}(X/R)[1]), \end{aligned}$$

and we have seen that $*$ acts trivially on $Q\mathcal{P}(A, 0)/G^1 = \mathcal{P}(A, 0)$, so $Q\mathcal{P}(A, 0)^{sd}/G^1 \simeq \mathcal{P}(A, 0)$. \square

Remark 2.21. The proof of Theorem 2.20 shows that for a self-dual quantisation of a non-degenerate 0-shifted Poisson structure, the w -compatible generalised symplectic structure is determined by its even coefficients. This raises the question of whether the odd coefficients must be homotopic to 0, as happens in the (-1) -shifted case by [Pri4, Remark 4.6]. The answer depends on the choice of w , as follows.

The involution i from Lemma 1.14 is not just a DGLA automorphism. If we write $f^t := -i(f)$, then $(f \cdot g)^t = (-1)^{\deg f \deg g} g^t \cdot f^t$ and $\{f\}\{g_1, \dots, g_m\}^t \simeq \mp \{f^t\}\{g_m^t, \dots, g_1^t\}$, so $(-)^t: C_R^\bullet(A)^{\text{opp}} \rightarrow C_R^\bullet(A)$ makes $C_R^\bullet(A)$ into an involutive brace algebra. The opposite brace algebra B^{opp} is most easily understood in terms of the associated B_∞ -algebra, which is a bialgebra structure on the tensor coalgebra $T(B[1])$: to form B^{opp} , we just take the opposite comultiplication on $T(B[1])$.

We can define an involution of the E_2 operad similarly, which takes an embedding $[1, k] \times I^2 \rightarrow I^2$ of k little squares in a big square, and reverses the order of the labels $[1, k]$ with appropriate signs. This involution comes from an element $t \in \text{GT}$ which maps to $-1 \in \mathbb{G}_m$; in other words, t is a (-1) -Drinfeld associator. It gives a notion of opposite E_2 -algebra, with $(-)^t: C_R^\bullet(A)^{\text{opp}} \rightarrow C_R^\bullet(A)$ then giving $C_R^\bullet(A)$ the structure of an involutive E_2 -algebra.

Levi decompositions w of GT with $w(-1) = t$ form a torsor $\text{Levi}_{\text{GT}}^t$ for the subgroup $(\text{GT}^1)^t$ of t -invariants in GT^1 , and for any such $w \in \text{Levi}_{\text{GT}}^t(\mathbb{Q})$, the ∞ -functor p_w sends opposite E_2 -algebras to opposite P_2 -algebras, defined by reversing the sign of the Lie bracket. This gives

$$\mu_w(\omega, \Delta)^t = \mu_w(\omega, -\Delta^t),$$

so $\omega(\hbar)$ is compatible with Δ if and only if $\omega(-\hbar)$ is compatible with Δ^* , implying that the odd coefficients of ω must be homotopic to 0 when Δ is non-degenerate and self-dual.

For a more explicit description of the generalised symplectic structure ω corresponding to a non-degenerate self-dual quantisation Δ , observe that we then have an isomorphism

$$\begin{aligned} \mu_w(-, \Delta): H^*(F^2\text{DR}(X/R) \times \hbar^2\text{DR}(X/R)[[\hbar^2]]) \\ \rightarrow \{v \in H^*(T_\Delta(\widehat{F^2 Q\text{Pol}}(A, 0))) : v(-\hbar) = v^t(\hbar)\}, \end{aligned}$$

and that $[\omega]$ must be the inverse image of $[\hbar^2 \frac{\partial \Delta}{\partial \hbar}]$.

Remark 2.22. Similarly to Remarks 2.9 and 2.19, we could consider the space $\mathbf{RLevi}_{\text{GT}}^t(R)$ of R -linear Levi decompositions with $w(-1) = t$, and the proof of Theorem 2.20 then combines with Remark 2.21 to give a canonical weak equivalence

$$\mathbf{RLevi}_{\text{GT}}^t(R) \times Q\mathcal{P}(A, 0)^{\text{nondeg}, sd} \simeq \mathbf{RLevi}_{\text{GT}}^t(R) \times \mathcal{P}(A, 0)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2\text{DR}(X/R)[[\hbar^2]][1]).$$

over $\mathbf{RLevi}_{\text{GT}}^t(R)$.

2.5. Comparison with Kontsevich–Tamarkin quantisations. In [Kon2], Kontsevich showed that for a smooth algebraic variety X over a field k , and a choice $w \in \text{Levi}_{\text{GT}}(k)$, every Poisson structure π lifts to an algebroid quantisation of X . We now investigate how this quantisation relates to our quantisations above when π is non-degenerate and $X = \text{Spec } A$ affine.

In Tamarkin's reinterpretation [Tam], the main intermediate step, as described in [Kon1] or [VdB, Theorem 1.1], is the existence of a canonical quasi-isomorphism

$$p_w \mathrm{CC}_k^\bullet(A) \simeq \prod_{i \geq 0} \mathrm{Hom}_A(\Omega_{A/R}^i, A)[-i]$$

of filtered $P_{2,\infty}$ -algebras, lifting the HKR isomorphism. This immediately gives us a $k[[\hbar]]$ -linear $P_{2,\infty}$ -algebra quasi-isomorphism

$$Q\widehat{\mathrm{Pol}}_w(A, 0) \simeq \widehat{\mathrm{Pol}}(A, 0)[[\hbar]]$$

sending the filtration $\{\tilde{F}^p\}_p$ on the left to $\{\prod_i F^{p-i} \hbar^i\}_p$.

In particular, we have a section ϕ of the projection $Q\widehat{\mathrm{Pol}}_w(A, 0) \rightarrow \widehat{\mathrm{Pol}}(A, 0)$, and the quantisation of [Kon2] is the induced map

$$\phi_w : \mathcal{P}(A, 0) \rightarrow Q\mathcal{P}(A, 0).$$

To understand how this interacts with our construction, note that functoriality of μ implies that $\mu_w(\omega, \phi_w(\pi)) = \phi_w(\mu(\omega, \pi))$, so $\phi_w(\pi)$ is w -compatible with a pre-symplectic form ω whenever π is compatible with ω .

In the non-degenerate setting $\omega = \pi^{-1}$, and in order to compare $\phi_w(\pi)$ with the quantisations of Theorem 2.20, we need to know whether it is self-dual. Tamarkin's approach to quantisation, as described in [Kon1], relies on showing that the ∞ -groupoid of invariant P_2 -algebra deformations of $\widehat{\mathrm{Pol}}(A, 0)$ is trivial. Morphisms being unique, the same will necessarily be true for involutive P_2 -algebra deformations. If $w(-1) = t$, this gives $\phi : \widehat{\mathrm{Pol}}(A, 0)[[\hbar^2]] \simeq Q\widehat{\mathrm{Pol}}_w(A, 0)^{sd}$, so we have shown the following:

Proposition 2.23. *For a smooth algebra A over a field $k \supset \mathbb{Q}$ and for $w \in \mathrm{Levi}_{\mathrm{GT}}^t(k)$, the Kontsevich–Tamarkin quantisation $\phi_w(\pi)$ of any Poisson structure π on A is self-dual. When π is non-degenerate, the quantisation $\phi_w(\pi)$ corresponds under Theorem 2.20 to the constant de Rham power series π^{-1} .*

Finally, observe that the explicit quantisation formulae of [Kon3] satisfy $a \star_{-\hbar} b = b \star_{\hbar} a$, so are self-dual and are thus covered by the proposition.

3. QUANTISATION FOR DERIVED STACKS

As in [Pri4, §3], in order to pass from bi-CDGAs to derived Artin stacks, we will exploit étale functoriality using Segal spaces.

3.1. Quantised polyvectors for diagrams.

Definition 3.1. Given a small category I , an I -diagram A in $DG^+ dg\mathrm{CAlg}(R)$, and an A -module M in I -diagrams of chain cochain complexes, define the filtered Hochschild cochain complex $\mathrm{CC}_R^\bullet(A, M)$ to be the equaliser of the obvious diagram

$$\prod_{i \in I} \mathrm{CC}_R^\bullet(A(i), M(i)) \rightrightarrows \prod_{f: i \rightarrow j \text{ in } I} \mathrm{CC}_R^\bullet(A(i), M(j)),$$

with the filtration $F_k \mathrm{CC}_R^\bullet(A, M)$ defined similarly, for the Hochschild complexes of Definition 1.12.

We then write $\mathrm{CC}_R^\bullet(A) := \mathrm{CC}_R^\bullet(A, A)$, which inherits the structure of a brace algebra from each $\mathrm{CC}_R^\bullet(A(i), A(i))$.

For $f: i \rightarrow j$ a morphism in I , observe that the HKR maps

$$\mathrm{gr}_k^F \mathrm{CC}_R^\bullet(A(i), f_* M(j)) \rightarrow \hat{\mathrm{Hom}}_{A(i)}(\Omega_{A(i)/R}^k, f_* M(j))$$

are quasi-isomorphisms whenever $A(i)$ is cofibrant in the model structure of Lemma 1.3. Also note that if $u: I \rightarrow J$ is a morphism of small categories and A is a functor from J to $DG^+ dg\mathrm{CAlg}(R)$ with $B = A \circ u$, then we have a natural map $\mathrm{CC}_R^\bullet(A) \rightarrow \mathrm{CC}_R^\bullet(B)$.

In order to ensure that $\mathrm{CC}_R^\bullet(A, M)$ has the correct homological properties, we now consider categories of the form $[m] = (0 \rightarrow 1 \rightarrow \dots \rightarrow m)$.

Lemma 3.2. *If A is an $[m]$ -diagram in $DG^+ dg\mathrm{CAlg}(R)$ which is cofibrant and fibrant for the injective model structure (i.e. each $A(i)$ is cofibrant in the model structure of Lemma 1.3 and the maps $A(i) \rightarrow A(i+1)$ are surjective), then $\mathrm{gr}_k^F \mathrm{CC}_R^\bullet(A)$ is a model for the derived Hom-complex $\mathbf{R}\hat{\mathrm{Hom}}_A(\mathbf{L}\Omega_A^k, A)$.*

Proof. The proof of [Pri5, Lemma 2.2] adapts verbatim to bi-CDGAs to give $\hat{\mathrm{Hom}}_A(\Omega_{A/R}^k, A) \simeq \mathbf{R}\hat{\mathrm{Hom}}_A(\mathbf{L}\Omega_A^k, A)$, from which this result follows immediately via the HKR isomorphism. \square

The constructions in §1 now all carry over verbatim, generalising from cofibrant bi-CDGAs to $[m]$ -diagrams of bi-CDGAs which are cofibrant and fibrant for the injective model structure. In order to identify $Q\mathcal{P}/G^1$ with \mathcal{P} , and for notions such as non-degeneracy to make sense, we have to assume that for our fibrant cofibrant $[m]$ -diagram A of bi-CDGAs, each $A(j)$ satisfies Assumption 1.7, so there exists N for which the chain complexes $(\Omega_{A(j)/R}^1 \otimes_{A(j)} A(j)^0)^i$ are acyclic for all $i > N$.

Definition 3.3. Given an $[m]$ -diagram A satisfying the conditions of Lemma 3.2, define

$$G\mathrm{PreSp}(A/R, 0) := G\mathrm{PreSp}(A(0)/R, 0) = \varprojlim_{i \in [m]} G\mathrm{PreSp}(A(i)/R, 0),$$

for the space $G\mathrm{PreSp}$ of generalised pre-symplectic structures of Definition 2.4.

Given a choice $w \in \mathrm{Levi}_{\mathrm{GT}}(\mathbb{Q})$ of Levi decomposition for GT, define

$$\mu_w: G\mathrm{PreSp}(A/R, 0) \times Q\mathcal{P}_w(A, 0) \rightarrow TQ\mathcal{P}_w(A, 0)$$

by setting $\mu_w(\omega, \Delta)(i) := \mu_w(\omega(i), \Delta(i)) \in TQ\mathcal{P}_w(A(i), 0)$ for $i \in [m]$, and let $Q\mathrm{Comp}_w(A, 0)$ be the homotopy vanishing locus of

$$(\mu_w - \sigma): G\mathrm{PreSp}(A/R, 0) \times Q\mathcal{P}_w(A, 0) \rightarrow TQ\mathcal{P}_w(A, 0).$$

over $Q\mathcal{P}_w(A, 0)$.

The following is [Pri5, Lemma 2.3]:

Lemma 3.4. *If $D = (A \rightarrow B)$ is a fibrant cofibrant $[1]$ -diagram in $DG^+ dg\mathrm{CAlg}(R)$ which is formally étale in the sense that the map*

$$\{\mathrm{Tot} \sigma^{\leq q}(\Omega_A^1 \otimes_A B^0)\}_q \rightarrow \{\mathrm{Tot} \sigma^{\leq q}(\Omega_B^1 \otimes_B B^0)\}_q$$

is a pro-quasi-isomorphism, then the map

$$\hat{\mathrm{Hom}}_D(\Omega_D^k, D) \rightarrow \hat{\mathrm{Hom}}_A(\Omega_A^k, A),$$

is a quasi-isomorphism for all k .

As in [Pri5, §3.4.2], for any of the constructions F based on $Q\mathcal{P}$, [Pri5, Definition 2.7] adapts to give an ∞ -functor

$$\mathbf{R}F: \mathbf{LDG}^+ dg\mathbf{CAlg}(R)^{\acute{e}t} \rightarrow \mathbf{LsSet}$$

with $(\mathbf{R}F)(A) \simeq F(A)$ for all cofibrant bi-CDGAs A over R , where $DG^+ dg\mathbf{CAlg}(R)^{\acute{e}t} \subset DG^+ dg\mathbf{CAlg}(R)$ is the subcategory of homotopy formally étale morphisms.

Immediate consequences of Propositions 2.16 and 2.17 and Theorem 2.20 are that for any $w \in \text{Levi}_{\text{GT}}(\mathbb{Q})$, the canonical maps

$$\begin{aligned} \mathbf{R}Q\text{Comp}_w(-, 0)^{\text{nondeg}} &\rightarrow \mathbf{R}Q\mathcal{P}_w(-, 0)^{\text{nondeg}} \simeq \mathbf{R}Q\mathcal{P}(-, 0)^{\text{nondeg}} \\ \mathbf{R}Q\mathcal{P}_w(-, 0)^{\text{nondeg}}/G^j &\rightarrow (\mathbf{R}Q\mathcal{P}_w(-, 0)^{\text{nondeg}}/G^2) \times_{\mathbf{R}(G\text{Sp}(-, 0)/G^2)}^h \mathbf{R}(G\text{Sp}(-, 0)/G^j) \\ \mathbf{R}Q\mathcal{P}_w(-, 0)^{\text{nondeg}, sd} &\simeq \mathbf{R}\mathcal{P}(-, 0)^{\text{nondeg}} \times \underline{\mathbf{MC}}(\hbar^2 \text{DR}(-/R)[[\hbar^2]][1]) \end{aligned}$$

are weak equivalences of ∞ -functors on the full subcategory of $\mathbf{LDG}^+ dg\mathbf{CAlg}(R)^{\acute{e}t}$ consisting of objects satisfying the conditions of Assumption 1.7, for all $j \geq 2$.

3.2. Hypergroupoids. We now recall the main constructions from [Pri3], as summarised in [Pri5, §2.2].

We require our chain CDGA R over \mathbb{Q} to be concentrated in non-negative chain degrees, and write $dg_+ \mathbf{CAlg}(R) \subset dg\mathbf{CAlg}(R)$ for the full subcategory of chain CDGAs which are concentrated in non-negative chain degrees. We denote the opposite category to $dg_+ \mathbf{CAlg}(R)$ by $DG^+ \text{Aff}_R$. Write $sDG^+ \text{Aff}_R$ for the category of simplicial diagrams in $DG^+ \text{Aff}_R$. A morphism in $DG^+ \text{Aff}_R$ is said to be a fibration if it is given by a cofibration in the opposite category $dg_+ \mathbf{CAlg}(R)$.

Definition 3.5. Given $Y \in sDG^+ \text{Aff}_R$, a DG Artin N -hypergroupoid X over Y is a morphism $X \rightarrow Y$ in $sDG^+ \text{Aff}_R$ for which:

- (1) the matching maps

$$X_m \rightarrow M_{\partial \Delta^m}(X) \times_{M_{\partial \Delta^m}(Y)} Y_m$$

are fibrations for all $m \geq 0$;

- (2) the partial matching maps

$$X_m \rightarrow M_{\Lambda_k^m}(X) \times_{M_{\Lambda_k^m}^h(Y)}^h Y_m$$

are smooth surjections for all $m \geq 1$ and k , and are weak equivalences for all $m > N$ and all k .

A morphism $X \rightarrow Y$ in $sDG^+ \text{Aff}_R$ is a trivial DG Artin (resp. DM) N -hypergroupoid if and only if the matching maps

$$X_m \rightarrow M_{\partial \Delta^m}(X) \times_{M_{\partial \Delta^m}(Y)} Y_m$$

are surjective smooth fibrations for all m , and are weak equivalences for all $m \geq n$.

The following is [Pri3, Theorem 4.15 and Corollary 6.35], as spelt out in [Pri2, Theorem 5.11]:

Theorem 3.6. *The ∞ -category of strongly quasi-compact N -geometric derived Artin stacks over R is given by localising the category of DG Artin N -hypergroupoids over R at the class of trivial relative DG Artin N -hypergroupoids.*

Given a DG Artin N -hypergroupoid X , we denote the associated N -geometric derived Artin stack by X^\sharp .

There is a denormalisation functor D from non-negatively graded CDGAs to cosimplicial algebras, with left adjoint D^* as in [Pri1, Definition 4.20]. Given a cosimplicial chain CDGA A , D^*A is then a bi-CDGA, with $(D^*A)_j^i = 0$ for $i < 0$.

3.3. Global quantisations. The following is [Pri5, Corollary 3.13], showing that a DG Artin N -hypergroupoid X can be recovered from the bi-CDGAs $D^*O(X^{\Delta^j})$:

Lemma 3.7. *For any simplicial presheaf F on $DGAff(R)$ and any Reedy fibrant simplicial derived affine X , there is a canonical weak equivalence*

$$\varprojlim_{j \in \Delta} \text{map}(\text{Spec } DD^*O(X^{\Delta^j}), F) \rightarrow \text{map}(X, F).$$

Lemma 3.7 and [Pri5, Proposition 3.18] ensure that if a morphism $X \rightarrow Y$ of DG Artin N -hypergroupoids becomes an equivalence on hypersheafifying, then $D^*O(Y) \rightarrow D^*O(X)$ is formally étale in the sense of Lemma 3.4. In particular this means that the maps $\partial^i: D^*O(X^{\Delta^j}) \rightarrow D^*O(X^{\Delta^{j+1}})$ and $\sigma^i: D^*O(X^{\Delta^{j+1}}) \rightarrow D^*O(X^{\Delta^j})$ are formally étale. Thus $D^*O(X^{\Delta^\bullet})$ can be thought of as a DM hypergroupoid in bi-CDGAs, and we may make the following definition:

Definition 3.8. Given a DG Artin N -hypergroupoid X over R and any of the constructions F based on $Q\mathcal{P}$, write

$$F(X) := \varprojlim_{j \in \Delta} \mathbf{R}F(D^*O(X^{\Delta^j})),$$

for $\mathbf{R}F$ as in §3.1.

The proof of [Pri5, Proposition 3.29] shows that if $Y \rightarrow X$ is a trivial DG Artin hypergroupoid, then the morphism $F(X) \rightarrow F(Y)$ is an equivalence for any of the constructions $F = \mathcal{P}, \text{Comp}, \text{PreSp}$. Thus the following is well-defined:

Definition 3.9. Given a strongly quasi-compact DG Artin N -stack \mathfrak{X} over R , define the spaces $Q\mathcal{P}(\mathfrak{X}/R, 0)$, $Q\text{Comp}_w(\mathfrak{X}/R, 0)$, $G\text{Sp}(\mathfrak{X}, 0)$ to be the spaces $Q\mathcal{P}(X/R, 0)$, $\text{Comp}_w(X/R, 0)$, $G\text{Sp}(X, 0)$ for any DG Artin N -hypergroupoid X with $X^\sharp \simeq \mathfrak{X}$.

Proposition 3.10. *For any strongly quasi-compact DG Artin N -stack \mathfrak{X} over R , there is a natural map from $Q\mathcal{P}(\mathfrak{X}, 0)$ to the space of curved A_∞ deformations $(\text{per}_{dg}(\mathfrak{X})[[\hbar]], \{m^{(i)}\}_{i \geq 0})$ of the dg category $\text{per}_{dg}(\mathfrak{X})$ of perfect $\mathcal{O}_{\mathfrak{X}}$ -complexes.*

This restricts to a map from $Q\mathcal{P}(\mathfrak{X}, 0)^{sd}$ to the space of involutive curved A_∞ deformations of $\text{per}_{dg}(\mathfrak{X})$.

Proof. Combining [Pri3, Proposition 5.12] with [Pri5, Lemma 3.8 and Corollary 3.13] and choosing a derived Artin N -hypergroupoid X representing \mathfrak{X} , we have

$$\text{per}_{dg}(\mathfrak{X}) \simeq \varprojlim_{j \in \Delta} \text{per}_{dg}(D^*O(X^{\Delta^j})).$$

By definition, $Q\mathcal{P}(\mathfrak{X}, 0) \simeq \varprojlim_{j \in \Delta} Q\mathcal{P}(D^*O(X^{\Delta^j}), 0)$, so the existence of the map follows from Proposition 1.23. The second statement is an immediate consequence. \square

Adapting [Pri4, Definition 2.21] to unshifted structures gives:

Definition 3.11. Given a Poisson structure $\pi \in \mathcal{P}(\mathfrak{X}, 0)$, we say that π is non-degenerate if the induced map

$$\pi^\sharp: \mathbf{L}\Omega_{\mathfrak{X}}^1 \rightarrow \mathbf{R}\mathcal{H}om_{\mathcal{O}_{\mathfrak{X}}}(\mathbf{L}\Omega_{\mathfrak{X}}^1, \mathcal{O}_{\mathfrak{X}})$$

is a quasi-isomorphism of sheaves on \mathfrak{X} , and $\mathbf{L}\Omega_{\mathfrak{X}}^1$ is perfect.

Combined with the results above, an immediate consequence of the generalisation of Propositions 2.16 and 2.17 and Theorem 2.20 in §3.1 is:

Theorem 3.12. *For any strongly quasi-compact DG Artin N -stack \mathfrak{X} over R , and any $w \in \text{Levi}_{\text{GT}}(\mathbb{Q})$, there are canonical weak equivalences*

$$\begin{aligned} Q\text{Comp}_w(\mathfrak{X}, 0)^{\text{nondeg}} &\rightarrow Q\mathcal{P}_w(\mathfrak{X}, 0)^{\text{nondeg}} \simeq Q\mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}} \\ Q\mathcal{P}_w(\mathfrak{X}, 0)^{\text{nondeg}}/G^j &\rightarrow (Q\mathcal{P}_w(\mathfrak{X}, 0)^{\text{nondeg}}/G^2) \times_{(G\text{Sp}(\mathfrak{X}, 0)/G^2)}^h (G\text{Sp}(\mathfrak{X}, 0)/G^j) \\ Q\mathcal{P}_w(\mathfrak{X}, 0)^{\text{nondeg}, sd} &\simeq \mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2\text{DR}(\mathfrak{X}/R)[[\hbar^2]][1]). \end{aligned}$$

Remarks 3.13. As in Remark 2.19, instead of choosing one Levi decomposition, we could work with the space $\mathbf{R}\text{Levi}_{\text{GT}}(R)$ of R -linear Levi decompositions, and the proof of Theorem 3.12 then gives equivalences

$$\begin{aligned} &\mathbf{R}\text{Levi}_{\text{GT}}(R) \times Q\mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}}/G^j \\ &\rightarrow \mathbf{R}\text{Levi}_{\text{GT}}(R) \times (Q\mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}}/G^2) \times_{(G\text{Sp}(\mathfrak{X}, 0)/G^2)}^h (G\text{Sp}(\mathfrak{X}, 0)/G^j), \\ &\mathbf{R}\text{Levi}_{\text{GT}}(R) \times Q\mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}, sd} \\ &\simeq \mathbf{R}\text{Levi}_{\text{GT}}(R) \times \mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2\text{DR}(\mathfrak{X}/R)[[\hbar^2]][1]) \end{aligned}$$

over $\mathbf{R}\text{Levi}_{\text{GT}}(R)$.

As in Remark 2.21, the power series w -compatible with a quantisation, although determined by its even coefficients, might have odd coefficients unless $w(-1) = t$. The reasoning of Remark 2.22 gives a canonical equivalence

$$\mathbf{R}\text{Levi}_{\text{GT}}^t(R) \times Q\mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}, sd} \simeq \mathbf{R}\text{Levi}_{\text{GT}}^t(R) \times \mathcal{P}(\mathfrak{X}, 0)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2\text{DR}(\mathfrak{X}/R)[[\hbar^2]][1])$$

over $\mathbf{R}\text{Levi}_{\text{GT}}^t(R)$ which does send a quantisation to its family of compatible de Rham power series.

As in Remark 1.32, self-dual quantisations of $\mathcal{O}_{\mathfrak{X}}$ also give rise to self-dual quantisations of all involutive \mathbb{G}_m -gerbes, and in particular of the Picard algebroid with involution given by $\mathbf{R}\mathcal{H}om_{\mathcal{O}_{\mathfrak{X}}}(-, \mathcal{L})$ for any line bundle \mathcal{L} .

Finally, applying étale descent to Proposition 2.23 shows that for a smooth DM stack X , the Kontsevich–Tamarkin quantisation $\phi_w(\pi)$ of any Poisson structure π on X is self-dual whenever $w \in \text{Levi}_{\text{GT}}^t(k)$. When π is non-degenerate, the quantisation $\phi_w(\pi)$ then corresponds under Theorem 3.12 to the constant de Rham power series π^{-1} .

Remark 3.14 (Positively shifted symplectic structures). All of the definitions of this paper adapt to n -shifted symplectic structures for $n > 0$, by working with E_{n+1} -Hochschild cohomology. Formality of E_{n+1} gives canonical P_{n+2} -algebra quasi-isomorphisms

$$Q\widehat{\text{Pol}}_w(A, n) \simeq \widehat{\text{Pol}}(A, n)[[\hbar]],$$

so existence of quantisations is automatic.

However, the methods of this paper adapt to yield a little more information for positively shifted symplectic structures, since they give equivalences

$$Q\mathcal{P}_w(\mathfrak{X}, n)^{\text{nondeg}, sd} \simeq \mathcal{P}(\mathfrak{X}, n)^{\text{nondeg}} \times \underline{\text{MC}}(\hbar^2\text{DR}(\mathfrak{X}/R)[[\hbar^2]][n+1])$$

for any $w \in \text{Levi}_{\text{GT}}^t$. This provides a complete parametrisation of non-degenerate self-dual E_{n+1} quantisations in terms of de Rham power series, with the choice of Levi decomposition only altering the parametrisation.

REFERENCES

- [Bra] Christopher Braun. Involutive A_∞ -algebras and dihedral cohomology. *J. Homotopy Relat. Struct.*, 9(2):317–337, 2014. arXiv:1209.1261v2 [math.QA].
- [CPT⁺] D. Calaque, T. Pantev, B. Toën, M. Vaquié, and G. Vezzosi. Shifted Poisson structures and deformation quantization. arXiv:1506.03699v3 [math.AG], 2015.
- [Hin] Vladimir Hinich. DG coalgebras as formal stacks. *J. Pure Appl. Algebra*, 162(2-3):209–250, 2001.
- [Kon1] Maxim Kontsevich. Operads and motives in deformation quantization. *Lett. Math. Phys.*, 48(1):35–72, 1999. Moshé Flato (1937–1998).
- [Kon2] Maxim Kontsevich. Deformation quantization of algebraic varieties. *Lett. Math. Phys.*, 56(3):271–294, 2001. EuroConférence Moshé Flato 2000, Part III (Dijon).
- [Kon3] Maxim Kontsevich. Deformation quantization of Poisson manifolds. *Lett. Math. Phys.*, 66(3):157–216, 2003.
- [LdB] Wendy Lowen and Michel Van den Bergh. The curvature problem for formal and infinitesimal deformations. arXiv:1505.03698, 2015.
- [Pet] Dan Petersen. Minimal models, GT-action and formality of the little disk operad. *Selecta Math. (N.S.)*, 20(3):817–822, 2014. arXiv:1303.1448v1 [math.AT].
- [Pri1] J. P. Pridham. Unifying derived deformation theories. *Adv. Math.*, 224(3):772–826, 2010. arXiv:0705.0344v5 [math.AG].
- [Pri2] J. P. Pridham. Notes characterising higher and derived stacks concretely. arXiv:1105.4853v3 [math.AG], 2011.
- [Pri3] J. P. Pridham. Presenting higher stacks as simplicial schemes. *Adv. Math.*, 238:184–245, 2013. arXiv:0905.4044v4 [math.AG].
- [Pri4] J. P. Pridham. Deformation quantisation for (-1) -shifted symplectic structures and vanishing cycles. arXiv:1508.07936v2 [math.AG], 2015.
- [Pri5] J. P. Pridham. Shifted Poisson and symplectic structures on derived N -stacks. arXiv:1504.01940v3 [math.AG], 2015.
- [Tam] Dmitry E. Tamarkin. *Operadic proof of M. Kontsevich’s formality theorem*. ProQuest LLC, Ann Arbor, MI, 1999. Thesis (Ph.D.)—The Pennsylvania State University.
- [VdB] Michel Van den Bergh. On global deformation quantization in the algebraic case. *J. Algebra*, 315(1):326–395, 2007.
- [Vor] Alexander A. Voronov. Homotopy Gerstenhaber algebras. In *Conférence Moshé Flato 1999, Vol. II (Dijon)*, volume 22 of *Math. Phys. Stud.*, pages 307–331. Kluwer Acad. Publ., Dordrecht, 2000. arXiv:9908040 [math.QA].